

Behavioral Risk Factor Surveillance by Race and Ethnicity, 2003

	Diabetes	Current Smoking	Overweight (BMI >=25 to 29.99)	Obese (BMI>=30)	Acute Alcohol Risk	No Health Insurance	Alcohol Risk
Bexar County Non Hispanic White	9.4%	21.2%	35.4%	25.7%	14.4%	6.4%	10.0%
Bexar County Hispanic	10.5%	19.1%	39.6%	27.4%	27.0%	30.9%	10.3%
Bexar County Total	10.4%	19.3%	37.9%	27.8%	19.6%	18.2%	9.7%
Texas Non Hispanic White	7.9%	23.7%	35.4%	23.3%	16.1%	15.1%	6.4%
Texas Hispanic	7.8%	18.7%	41.1%	25.9%	19.4%	50.4%	5.7%
Texas Black	10.5%	20.4%	37.8%	34.9%	11.5%	26.8%	4.4%
Texas Total	8.1%	22.1%	36.9%	24.6%	16.3%	26.7%	5.9%

Bexar County 2003 Behavioral Risk Factor Surveillance by Race and Ethnicity



Texas 2003 Behavioral Risk Factor Surveillance by Race and Ethnicity



Note: 2003 Behavioral Risk Factor Surveillance Survey (BRFSS) included 329 randomly selected Bexar county residents.

Note: There were insufficient responses from the Black population in Bexar County to reliably estimate risk factors for 2003.

Note: Acute alcohol risk = 5 or more drinks on an occasion in the past 30 days.

Note: Heavy alcohol risk = > 2 drinks per day men, > 1 drink per day women - during the past 30 days.

 $\label{eq:Note:All reported rates (\%) are weighted for Texas demographics and the probability of selection and$

thus are not derived from the simple division of numerator and denominator cases.

Source: Behavioral Risk Factor Surveillance System, 2003.

Prepared by: Community Assessment Team

Center for Health Statistics

Texas Department of Health

June 23, 2004